REFERENCES

Arambewela L.S.R.,Arawwawala L.D.A.M.,Ratnasoorya W.D.,(2005).Antidiabetic activity of aqueous and ethanolic extract of *Piper betle* leaves in rats. Journal of ethnopharmacology,102:239-245.

Ahmed O.A.Z.,Safa Y S.,Assaf H.M.,Randa H.A.H.,(2005).Antidiabetic activity and toxicity of *Zizyphus spina-christi* leaves. Journal of ethnopharmacology,101:129-138.

Arulmozhi S.,Mazumder P.M.,Lohidasan S.,Thakurdesai P.,(2010). Antidiabetic and antihyperlipidemic activity of leaves of Alstonia scholaris Linn. European Journal of Integrative Medicine,2:23–32.

Are P.C.,Reddy R.R.,Adidala.,Puchchakayala G.,(2011). Hypoglycemic and Antidiabetic Activity of Glochidion velutinum on Streptozotocin-Nicotinamide Induced Type 2 Diabetic Rats. European Journal of Biological Sciences,3(4): 126-130.

Bhat, M., Zingarde, S.S., Bhargava, S.Y., Kumar, A.R., Joshi, B.N., (2008). Antidiabetic Indian plants: a good source of potent amylase inhibitors. Evidence Based on Complementary and Alternative Medicine, doi:10.1093/ecam/nen040.

Batra S, Srivastava S, Singh K, Chander C, Khanna AK.,(2000). Synthesis & biological evaluation of 3-substituted amino-1-aryl-6-hydroxy-hex-2-ene-1-one as antioxidant & hypolipidemic agents. Bioorganic and med chem. 8:2195-2209.

Chakrabarti R., Reeba K.et al.,(2002). Antidiabetic and hypolipidemic activity of Helicteres isora in animal models. Journal of Ethnopharmacology,81:343-349.

Eidi A.,Eidi M.,(2009). Antidiabetic effects of sage (Salvia officinalis L.) leaves in normal and streptozotocin-induced diabetic rats. Diabetes & Metabolic Syndrome: Clinical Research & Reviews,3:40–44.

Eliza J.,Daisy P., Ignacimuthu S.,Duraipandiyan V.,(2009). Antidiabetic and antilipidemic effect of eremanthin from Costus speciosus (Koen.)Sm., in STZ-induced diabetic rats. Chemico-Biological Interactions,182:67–72.

Gaur K.,Rana A.C.,Nema R.K.,Kori M.L.,Sharma C.S.,(2009). Anti-inflammatory and analgesic activity of hydro-alcoholic leaves extract of euphorbia neriifolia linn. Asian Journal of Pharmaceutical and Clinical Research,2(1):26-29.

Jain S., Bhatia G.,Barik R.,Kumar P.,Jain A.,Dixit V.K.,(2010). Antidiabetic activity of Paspalum scrobiculatum Linn. in alloxan induced diabetic rats. Journal of Ethnopharmacology,127:325–328.

Kannur D.M.,Hukkeri V I.,Akki K.S.,(2006). Antidiabetic activity of Caesalpinia bonducella seed extracts in rats. Fitoterapia,77:546-549.

Kasiappan R.et al.,(2005). Antihyperlipidemic effect of *Eugenia jambolana* seed kernel on streptozotocin-induced diabetes in rats.Food and chemical toxicology,43(9):1433-1439.

KehkashanP., SiddiquiW.A.,(2011).Protective effect of butea monosperma on high-fat di et and streptozotocin-induced non-genetic rat model of type 2 diabetes: biochemical and histological evidences. International Journal of Pharmacy and Pharmaceutical Sciences,3(3):74-81.

Maiti R.,Jana D.,Das U.K.,Ghosh D.,(2004). Antidiabetic effect of aqueous extract of seed of Tamarindus indica in streptozotocin-induced diabetic rats. Journal of Ethnopharmacology,92:85–91.

Mohammed H., Daghriri H.A.et al.,(2008). Antidiabetic effect of alcoholic extract of Caralluma sinaica L. on streptozotocin-induced diabetic rabbits. Journal of Ethnopharmacology,117:215–220.

Nagarajan N.S.,Thirupathy P.K.,Radhad N.,Muralid A.,(2005). Antidiabetic and antihyperlipemic effects of Clemeo feline. Fitoterapia,76:310–315.

Nagappa A.N.,Thakurdesai P.A.,Rao N.V.,Singh J.,(2003). Antidiabetic activity of Terminalia catappa Linn fruits. Journal of Ethnopharmacology,88:45–50.

Ndiaye M.et al.,(2008). Antidiabetic properties of aqueous barks extract of Parinari excelsa in alloxan-induced diabetic rats. Fitoterapia,79:267 – 270.

Oh W.K.et al.,(2005). Antidiabetic effects of extracts from Psidium guajava. Journal of Ethnopharmacology,96:411–415.

Pari L.,Amarnath S.,(2004). Antidiabetic activity of Boerhaavia diffusa L. effect on hepatic key enzymes in experimental diabetes. Journal of ethnopharmacology,91:109-113.

Rao B.K.,Sudarshan P.R., Rajasekhar M.D.,Nagaraju N.,Rao A.,(2003). Antidiabetic activity of Terminalia pallida fruit in alloxan induced diabetic rats. Journal of Ethnopharmacology,85:169–172.

Rao B.K.,Kesavulu M.M.,Giri R., Rao A.,(1999). Antidiabetic and hypolipidemic effects of Momordica cymbalaria Hook. fruit powder in alloxan-diabetic rats. Journal of Ethnopharmacology,67:103–109.

Raut N.A., Gaikwad N.J.,(2006). Antidiabetic activity of hydro-ethanolic extract of Cyperus rotundus in alloxan induced diabetes in rats. Fitoterapia,77:585 – 588.

Sharma V.,Janmeda P.,Singh L.,(2011).A review on *Euphorbia neriifolia*(Sehund).Sputula DD.1(2):107-111.

Sharma V.et al.,(2011). Chemoprotective activity of hydro-ethanolic extract of Euphorbia neriifolia Linn leaves against DENA-induced liver carcinogenesis in mice. Biology and Medicine,3 (2) : 36-44.

Sharma V.,Pracheta.,Paliwal R.,Sharma S.,(2011). Preliminary Phytochemical Screening and in vitro Antioxidant Potential of Hydro-Ethanolic extract of Euphorbia neriifolia Linn. International Journal of PharmTech Research,3:124-132.

SharmaV.,Pracheta.,PaliwalR.,SharmaS.,(2011).in vitro free radical scavenging and antio xidant potential of ethanolic extract of euphorbia neriifolia linn. International Journal of Pharmacy and Pharmaceutical Sciences, 3(1):238-242.

Sharma M.et al.,(2011). Evaluation of Antidiabetic and Antioxidant Effects of Seabuckthorn (Hippophae rhamnoides L.) in Streptozotocin-Nicotinamide Induced Diabetic Rats. The Open Conference Proceedings Journal,2:53-58.

Shabeer J.,Srivastava R.S.,Singh S.K.,(2009). Antidiabetic and antioxidant effect of various fractions of Phyllanthus simplex in alloxan diabetic rats. Journal of Ethnopharmacology,124:34–38.

Shokeen P.,Anand P.Murali K.Y.,Tandon V.,(2008). Antidiabetic activity of 50% ethanolic extract of Ricinus communis and its purified fractions. Food and Chemical Toxicology ,46 :3458–3466.

Singh R.K.,Mehta S.,Jaiswal D.,Rai P.K.,Watal G.,(2009). Antidiabetic effect of Ficus bengalensis aerial roots in experimental animals. Journal of Ethnopharmacology,123:110–114.

Sunila C.,Ignacimuthu S.,Agastian P.,(2011). Antidiabetic effect of Symplocos cochinchinensis (Lour.) S. Moore. in type 2 diabetic rats. Journal of Ethnopharmacology, 134:298–304.

Somani R.,Kasture S.,Singhai A.K.,(2006). Antidiabetic potential of Butea monosperma in rats. Fitoterapia,77:86–90.

Suba V.et al.,(2004). Antidiabetic potential of Barleria lupulina extract in rats. Fitoterapia, 75:1–4.

Shirwaika A.,Rajendran K.,Barik R.,(2006). Effect of aqueous bark extract of Garuga pinnata Roxb. in streptozotocin-nicotinamide induced type-II diabetes mellitus. Journal of Ethnopharmacology,107:285–290.

Srinivasan K.,Viswanad B.,Asrat L.,Kaul C.L.,Ramarao P.,(2005). Combination of high-fat diet-fed and low-dose streptozotocin-treated rat: A model for type 2 diabetes and pharmacological screening. Pharmacological Research 52:313–320.

Swamy M K.,Pokharen N.,Dahal S.,Anuradha M.,(2011). Phytochemical and antimicrobial studies of leaf extract of Euphorbia neriifolia. Journal of Medicinal Plants Research,5(24):5785-5788.

World Health Organization (WHO), (2006). Diabetes Programme. Available at http://www.who.Int/diabetes/en/ (accessed 15.10.07.).